

**Table III: Results of TPA Thrombolysis for Arterial Thromboses in Newborn Infants**

Author	N	Dose mg/kg/hr	Duration hr	Total Dose mg/kg	Heparin Concomitant	Lysis			Bleeding	Outcome
						C	P	N		
Levy, 1991 <sup>12</sup>	2	0.46; 0.47	2hr-3 days	1-34	No	2			1 site	Recovered
Dillon, 1992 <sup>13</sup>	1	0.5	Bolus	0.5	No	1			1 site	Recovered
Zenz, 1993 <sup>14</sup>	2	0.5 1 <sup>st</sup> hr 0.25 subsequent	4 hrs; 9 hrs	1.25; 2.5	No	1	1		None	Recovered
Farnoux, 1997 <sup>16</sup>	4	0.1 bolus; 0.3 infusions	1-3 three hr	1-3	No	3	1		None	Kidney lost in 2
Weiner, 1998 <sup>20</sup>	7	0.1-0.5	6-39 hrs	0.6-15.6		4	2	1	2 ICH (+ Pulm in 1) 3 minor	2 deaths
Knöfler, 2001 <sup>17</sup>	1	0.04	96	3.84	Yes	1			None	Recovered
Wang, 2001 <sup>18</sup>	3	0.1; 0.15; 0.5	3, 15, 16 hrs	0.3-8	Yes	3			3 at site 1 epistaxis 1 hematuria, hematoma	Recovered