## **UNC Health Science Initial Immunization Record**

Student Name: ID#: Tetanus/Diphtheria/Pertussis - Students must have at least 3 doses; one of which must be a Tdap booster and one of which must be within the past 10 yrs. (#1) mo./day/year (#2) mo./day/year (#3) mo./day/year DTP. DTaP or Td Tdap booster \*\*Must have one documented Measles/Mumps/Rubella - 2 doses of MMR at least 28 days apart after 12 months of age OR 2 doses of Measles and 2 doses of Mumps at least 28 days apart after 12 months of age and 1 dose of Rubella after 12 months of age OR laboratory proof of immunity (blood titer) to measles/mumps/rubella. If titers are negative or equivocal, repeat MMR series with doses at least 28 days apart. No titer is required after series repeat. (#1) mo./day/year (#2) mo./day/year MMR - 2 required on or after 1st birthday OR Measles – 2 required on or after first birthday Mumps – 2 required on or after first birthday Rubella - 1 required on or after first birthday OR Date of Titer Result MMR Titer \*must attach laboratory results Varicella – 2 doses of Varicella at least 4 weeks apart OR laboratory proof of immunity to varicella. If titer is negative or equivocal, repeat Varicella series with doses at least 4 weeks apart. No titer is required after series repeat. (#1) mo./day/year (#2) mo./day/year Varicella - 2 doses OR Date of Titer Result Varicella Titer \*must attach laboratory results Hepatitis B – a primary series of Hepatitis B vaccines AND a positive (>10 mIU/mL) serological quantitative Hepatitis B surface antibody titer (HBsAb) 1-2 months after the date of the last vaccine. If series was completed in the remote past, and if the titer checked upon matriculation is negative, student will get 1 hepatitis B vaccine dose and re-titer at least 1-2 months after vaccine. If the second titer is negative, student will get the additional Hepatitis B vaccine(s) to complete the series per the standard schedule. A final titer should be done 1-2 months after the final vaccine and if this is negative, the student should be considered a non-responder and evaluated and counseled appropriately. Those students recently vaccinated with a negative titer after a primary series can receive a second series with a re-titer 1-2 months after the final dose. Non-responders should be counseled and evaluated appropriately. (#1) mo./day/year (#2) mo./day/year (#3) mo./day/year Hepatitis B Series — a primary series required Date of Titer Result Hepatitis B Quantitative Titer \*must attach laboratory results Hepatitis B Series Repeat Date of Titer Result Hepatitis B Quantitative Titer Repeat \*must attach laboratory results Tuberculin Screening – IGRA Blood Test (preferred) OR a 2-step TB skin test (TST) placed within the past 12 months. 1st TST Place date 1st TST Read Date 2<sup>nd</sup> TST Place Date 2<sup>nd</sup> TST Read date 2 Step TST – placed within the past 12 months, must have been performed in the United States. The 2nd TST must be placed at least 1 week AFTER the 1st TST read date. OR Date of IGRA Result IGRA TB Screening \*must attach laboratory results \_\_ T-spot Quantiferon Gold Influenza – within the current season (AUG – DEC) Influenza Vaccine \*must attach laboratory results Health Care Provider's Signature AND Stamp Date